SLURP1 gene

secreted LY6/PLAUR domain containing 1

Normal Function

The *SLURP1* gene provides instructions for making a protein called secreted Ly6/ uPAR-related protein-1 (SLURP-1). This protein is found in skin cells and other cells that line the surfaces and cavities of the body. Like other Ly6/uPAR-related proteins, SLURP-1 folds into a particular shape and is thought to attach (bind) to other proteins called receptors to carry out signaling within cells. However, SLURP-1's role in the skin and the rest of the body is not completely understood.

Laboratory studies show that SLURP-1 can bind to nicotinic acetylcholine receptors (nAChRs). SLURP-1 specifically interacts with the alpha7 (α 7) subunit, which is a piece of some nAChRs. Nicotinic acetylcholine receptors are best known for their role in chemical signaling between nerve cells, but they are also found in other tissues. In the skin, nAChRs regulate the activity of genes involved in the growth and division (proliferation), maturation (differentiation), and survival of cells. Through its interaction with these receptors, SLURP-1 may be involved in skin growth and development.

Health Conditions Related to Genetic Changes


mal de Meleda

At least 15 mutations in the *SLURP1* gene have been found to cause mal de Meleda, a rare disorder characterized by tough, thickened skin on the hands and feet. On the palms and soles, the thickening is known as palmoplantar keratoderma; the thickened skin also extends to the backs of the hands and feet and up to the wrists and ankles. The *SLURP1* gene mutations involved in this condition lead to production of an altered SLURP-1 protein that is unstable and quickly broken down, if any protein is produced at all. As a result, affected individuals have little or no SLURP-1 protein. It is unclear how a lack of this protein leads to the skin problems that occur in mal de Meleda. Researchers speculate that without SLURP-1, the activity of genes controlled by nAChR signaling is altered, leading to overgrowth of skin cells or survival of cells that normally would have died. The excess of cells can result in skin thickening. It is unclear why skin on the hands and feet is particularly affected.

Chromosomal Location

Cytogenetic Location: 8q24.3, which is the long (q) arm of chromosome 8 at position 24.3

Molecular Location: base pairs 142,740,944 to 142,742,411 on chromosome 8 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- anti-neoplastic urinary protein
- ANUP
- ARS
- ARS(component B)-81/S
- ArsB
- LY6LS
- lymphocyte antigen 6-like secreted
- MDM
- secreted Ly-6/uPAR-related protein 1
- secreted Ly-6/uPAR-related protein 1 precursor
- secreted Ly6/uPAR related protein 1

Additional Information & Resources

Educational Resources

 Basic Neurochemistry: Molecular, Cellular and Medical Aspects (sixth edition, 1999): Nicotinic Receptors https://www.ncbi.nlm.nih.gov/books/NBK28261/

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28SLURP1%5BTIAB%5D%29+OR+%28Ly6/uPAR-related+protein+1%29+OR+%28ARS+B%29+OR+%28SLURP-1%29+OR+%28ARS+component+B%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 SECRETED LY6/PLAUR DOMAIN-CONTAINING PROTEIN 1 http://omim.org/entry/606119

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/GC SLURP1.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=SLURP1%5Bgene%5D
- HGNC Gene Family: LY6/PLAUR domain containing http://www.genenames.org/cgi-bin/genefamilies/set/1226
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=18746
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/57152
- UniProt http://www.uniprot.org/uniprot/P55000

Sources for This Summary

- Arredondo J, Chernyavsky AI, Webber RJ, Grando SA. Biological effects of SLURP-1 on human keratinocytes. J Invest Dermatol. 2005 Dec;125(6):1236-41.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16354194
- Chimienti F, Hogg RC, Plantard L, Lehmann C, Brakch N, Fischer J, Huber M, Bertrand D, Hohl D. Identification of SLURP-1 as an epidermal neuromodulator explains the clinical phenotype of Mal de Meleda. Hum Mol Genet. 2003 Nov 15;12(22):3017-24. Epub 2003 Sep 23.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/14506129

- Eckl KM, Stevens HP, Lestringant GG, Westenberger-Treumann M, Traupe H, Hinz B, Frossard PM, Stadler R, Leigh IM, Nürnberg P, Reis A, Hennies HC. Mal de Meleda (MDM) caused by mutations in the gene for SLURP-1 in patients from Germany, Turkey, Palestine, and the United Arab Emirates. Hum Genet. 2003 Jan;112(1):50-6. Epub 2002 Oct 19.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/12483299
- Favre B, Plantard L, Aeschbach L, Brakch N, Christen-Zaech S, de Viragh PA, Sergeant A, Huber M, Hohl D. SLURP1 is a late marker of epidermal differentiation and is absent in Mal de Meleda. J Invest Dermatol. 2007 Feb;127(2):301-8. Epub 2006 Sep 28.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/17008884
- Fischer J, Bouadjar B, Heilig R, Huber M, Lefèvre C, Jobard F, Macari F, Bakija-Konsuo A, Ait-Belkacem F, Weissenbach J, Lathrop M, Hohl D, Prud'homme JF. Mutations in the gene encoding SLURP-1 in Mal de Meleda. Hum Mol Genet. 2001 Apr 1;10(8):875-80. *Citation on PubMed:* https://www.ncbi.nlm.nih.gov/pubmed/11285253
- OMIM: SECRETED LY6/PLAUR DOMAIN-CONTAINING PROTEIN 1 http://omim.org/entry/606119

Reprinted from Genetics Home Reference: https://ghr.nlm.nih.gov/gene/SLURP1

Reviewed: November 2014 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services